



Wichita State University Libraries
SOAR: Shocker Open Access Repository

Francis Leland Russell

Biological Sciences

Phenological synchrony affects interaction strength of an exotic weevil with Platte thistle, a native host plant

F. Leland Russell

University of Nebraska-Lincoln, leland.russell@wichita.edu

Svata M. Louda

University of Nebraska-Lincoln

Recommended citation

Russell, F. Leland and Svata M. Louda. 2004. Phenological synchrony affects interaction strength of an exotic weevil with Platte thistle, a native host plant. *Oecologia* 139: 525–534.

This paper is posted in Shocker Open Access Repository

<http://soar.wichita.edu/dspace/handle/10057/3406>

**Phenological Synchrony Affects Interaction Strength of an Exotic
Weevil with Platte Thistle, a Native Host Plant**

F. Leland Russell and Svata M. Louda

School of Biological Sciences

University of Nebraska-Lincoln

Lincoln NE 68588-0118 USA

Corresponding Author:

F. Leland Russell

School of Biological Sciences

University of Nebraska-Lincoln

Lincoln NE 68588-0118

EMAIL: FRussell@unlnotes.unl.edu, FAX: 402-472-2083, TELEPHONE: 402-472-1539

Second author (Svata M. Louda) - EMAIL: Slouda@unl.edu, TELEPHONE: 402-472-

2763

Abstract

Phenological synchrony of a consumer population with its resource populations is expected to affect interaction intensity. We quantified phenological variation and synchrony of populations of an invasive Eurasian flower head weevil, *Rhinocyllus conicus*, that consumes florets, ovules, and seeds of developing flower heads of a native North American thistle, *Cirsium canescens*, in Sand Hills prairie in Nebraska USA. Variation in timing of adult activity among weevil populations was larger than variation in timing of flower head development among *C. canescens* populations, and it drove the observed variation in the phenological synchrony between weevil and host plant populations. Furthermore, the degree of phenological synchrony between populations was significant in explaining variation in weevil egg load on the newly acquired hostplant. Because population growth of *C. canescens* is limited by pre-dispersal seed losses to floral herbivores, variation in the synchrony of herbivore and plant flowering will affect the density of the plant population. These results provide strong quantitative support for the hypothesis that the synchrony of insect activity with plant resources can determine the magnitude of impact of floral herbivores on their host plant populations.

Key Words. Floral herbivory, invasive species, biocontrol, predispersal seed predation, plant population dynamics

Introduction

Temporal and spatial variation in the seasonal phenology of insects and their host plants represent important parameters in the ecology and evolution of herbivore-plant interactions (Augspurger 1981; Hunter 1992; Stenseth and Mysterud 2002). Masting, the between-year variation in seed resource availability, and its consequences for consumer-plant interactions has been examined intensively (Elkinton et al. 1996; Herrera et al. 1998; Schnurr et al. 2002). However, the number of studies that have examined the consequences of temporal and spatial variation in within-season phenological synchrony, the relative timing of consumer and resource populations (Rathcke and Lacey 1985), for populations of the resource species is much smaller.

Studies of within-season phenological synchrony have focused on two main issues. For plants, the main issue has been how seasonal variation in timing of the interacting species affects the level of herbivory on individual plants and individual plant fitness. These studies typically examine variation among individuals in: (1) flowering or leaf expansion (Mopper and Simberloff 1995; Pilson 2000), (2) flowering synchrony with others in the population (Augspurger 1981; Ollerton and Lack 1998), or (3) flowering strategy, i.e., synchronous vs. gradual flower presentation (Eriksson 1995) (Table 1: Questions 1-3). We found no studies that addressed the consequences of variation in the relative timing of insects and their host plants for host plant populations. For insects, the focus has been on effects of seasonal variation in the phenological synchrony of insects and plants on insect population dynamics (Table 1: Questions 5-6). These studies often address the relationship between individual plant phenology and insect abundance on the plant (Crawley and Akhteruzzaman 1988; Hodkinson et al. 2001; Hunter 1992), or the

role of insect-plant synchrony in insect outbreaks (Hunter and Elkinton 2000; Kerslake and Hartley 1997).

Unexpectedly, in a voluminous literature on the seasonal phenology of insect-plant interactions, we found only 18 studies that quantified the within-season synchrony of populations of both a consumer and its resource (Table 2). More studies of the effects of phenological mis-matches on consumer and resource population dynamics are needed (Stenseth and Mysterud's 2002), especially because climate change may differentially affect the phenologies of species at different trophic levels (Buse and Good 1996; Inouye et al. 2000; Visser and Holleman 2001). Furthermore among these 18 studies, we found that measures of phenological synchrony of insect and plant populations were inconsistent. Yet, we found no evaluation or discussion of the relative merits of the most commonly used measures of phenological synchrony between populations.

In this study, we evaluated the degree of within-season phenological synchrony between populations of an invasive biocontrol weevil, *Rhinocyllus conicus*, a Eurasian thistle flower head feeder, and the floral resources presented by populations of a newly adopted, native North American host plant, Platte thistle (*Cirsium canescens* Nutt.). Synchrony between flower head development by *C. canescens* and oviposition activity of *R. conicus* adults was hypothesized to explain the extent of *R. conicus* herbivory on this host species in the Sand Hills mid-grass prairie of the central Great Plains (Louda 1998). We used a long-term (12 yr) data set on Platte thistle flowering, augmented by 4 yrs of intensive measurement of weevil activity patterns, to ask specifically: Does variation in the phenological synchrony of *R. conicus* and floral resources of *C. canescens* explain variation in *R. conicus* oviposition and damage among *C. canescens* populations, within

or among years? We also examined the role of key climatic variables in explaining temporal and spatial variation in reproductive phenology of *C. canescens* populations. Finally, we reviewed the strengths and constraints of the various measures used to quantify the seasonal synchrony of consumer and resource populations.

Methods

Sites and Natural History

Field data were collected at eight sites in two mid-grass prairie preserves owned by The Nature Conservancy in the Nebraska Sand Hills, in the upper Great Plains USA. The two preserves were: Arapaho Prairie Preserve (Arthur County) in the southwest and Niobrara Valley Preserve (Brown County) in the northcentral Sand Hills, 270 km from Arapaho Prairie.

The flower head weevil, *Rhinocyllus conicus*, is native to Europe and was introduced deliberately into North America in 1969 against weedy exotic thistles, especially musk (nodding) thistle, *Carduus nutans* L. (Zwölfer and Harris 1984). The weevil invaded the two Sand Hills preserves in 1992-1993, far from the tallgrass habitat where musk thistle occurs (Louda 1998; Louda et al. 1997). Weevils oviposit on thistle flower heads that have not yet exerted florets. Eggs are laid under externally obvious egg cases made of masticated plant tissues. Larvae develop in the flower heads, consuming receptacle tissues, florets, ovules and seeds of the developing inflorescences. Development takes 53-76 d, allowing one generation per year (Zwölfer and Harris 1984).

Platte thistle (*Cirsium canescens* Nutt.) is restricted to the northcentral Great Plains, occurring in prairies on sand and gravel soils; the center of its distribution is the

Nebraska Sand Hills (Great Plains Flora Association 1986). Platte thistle is a tap-rooted, relatively short-lived, monocarpic perennial. The earliest flower heads are initiated in early May and flower in late May to mid June. Flower head development is determinant, and flower head position is a good index of relative flower head timing of development.

Flower head development and seed maturation are completed by the end of June.

Densities in Platte thistle populations are limited by seed input (Louda and Potvin 1995b). Therefore, additional ovule and seed consumption by *R. conicus* larvae likely is reducing densities of *C. canescens* populations further (Louda and Arnett 2000).

Data Collection

Flowering and floral herbivory by all insects on *C. canescens* were quantified by sampling adult plants at each preserve 1990-2001. In 1997-2000 we added intensive, twice weekly measurements of insect numbers on plants in at least two sites per preserve (Niobrara: inner and outer Salzman 1997-2000, and Mahoney 1998-2000; Arapaho: at least two of the following, JBO, RBO, West Foothills or Wilson, depending on the availability of flowering plants). At each site in each year (1990-2001), we marked every bolting *C. canescens* plant located along walking transects in late May ($12 \geq N \geq 5$). Lower sample sizes in some sites and years reflect low densities of bolting *C. canescens*. Platte thistle is a relatively sparse plant species; on average, densities of bolting Platte thistles are < 0.1 plant / hectare across the Sand Hills (Louda, unpublished data)

Flowering plants develop rapidly in May and mature seed in June; we measured plant development and all flower heads on each plant three (or four) times during this flowering period: (1) very early (early May) in most years, at the beginning of flower head initiation; (2) early-mid season (late May), when *R. conicus* adult activity and *C.*

canescens flower head initiation were at peak levels; and (3) mid-late season (late June), when *R. conicus* had finished oviposition and *C. canescens* had finished maturing most of its seed. We verified that we had not missed any later plant flower development by sampling again (4) late season (mid-late July), when no adult *R. conicus* remained and any late flower heads had matured. On each date, plant height, flower head development, and size (diameter, mm) of all heads were recorded. Development categories for individual flower heads were; “1”- no florets exerted and diameter < 15 mm, “2”-no florets and diameter \geq 15 mm, “3”- > 1 floret exerted, “4” - full flowering, “5” - post-flowering (following Lamp 1980).

We counted all insects present, including *R. conicus* egg cases per head after invasion, on each sampling date 1990-2001. For 1997-2000, to better quantify within-season phenology of *R. conicus* abundance and oviposition, we also counted insects at 4 d intervals (range 3–5 d) May – July, and noted number of flower heads available.

Daily temperature and precipitation data 1990-2001 were obtained from automated weather network stations at Arapaho and Ainsworth NE, 20 km south of Niobrara.

Statistical Analyses

Weevil Oviposition. We used Kruskal-Wallis tests (Sokal and Rohlf 1995) to compare *R. conicus* oviposition on *C. canescens* among years for each preserve, and a Wilcoxon ranked sum test to compare oviposition between preserves. Oviposition use by *R. conicus* was quantified for each plant as the mean number of egg cases per flower head initiated for each plant. We divided the number of egg cases on individual plants by the

number of heads initiated by the plant to take into account that larger plants could have more egg cases because they produced more flower heads.

Phenology and Synchrony. To quantify weevil phenology for preserves and sites, we calculated the proportion of the total cumulative number of weevils observed during a growing season that were seen by late May when oviposition activity was highest. The cumulative number of weevils observed at a site or preserve during a growing season was the sum of the weevil counts made at 4 d intervals (1997 - 2000). If weevils were observed on our first visit, we corrected the cumulative total by using the rate of increase in weevil numbers for the years when measurements started before weevils appeared. To quantify flower head development phenology of *C. canescens* plants for preserves and sites, we calculated the proportion of the total number of flower heads produced per plant over the season (1990-2001) that were initiated by late May when flower head development was peaking. We quantified the degree to which *R. conicus* phenology was ahead of *C. canescens* phenology as the proportion of adult weevils observed minus the mean proportion of *C. canescens* flower heads initiated by late May. We refer to this difference as the “degree *R. conicus* preceded *C. canescens*.” This measure was positive when *R. conicus* was earlier than *C. canescens*, and negative when *R. conicus* was later than *C. canescens*. The absolute value of this measure indicates the magnitude of the asynchrony of *R. conicus* and *C. canescens* populations. Larger absolute values indicate greater asynchrony than smaller absolute values.

We quantified phenologies of *R. conicus* and *C. canescens* populations in late May because our previous data indicated that, on average, both weevil activity and plant floral development were peaking at this time (Louda 1998; Louda 1999). We used the late May

sampling date as the criterion for investigation of relative phenologies based on the biology of the interaction; this was the time during which we could identify maximum differences in weevil activity and flower head initiation. The data from June and July were required to tally cumulative total number of eggs laid and cumulative flower heads produced, but were not informative on variation in the development of the interaction.

We used a two-way ANOVA to test for significant variation in *C. canescens* phenology among preserves and years, using Fischer's LSD to evaluate differences. Proportions were arcsine-transformed. We used a 2 x 4 x 2 contingency table analysis to determine whether the number of adult weevils observed by late May varied significantly among preserves and years, controlling for among preserve and year differences in the total number of adult weevils observed. To conduct *a posteriori* comparisons among preserves in each year, we used 2 x 2 contingency table analyses, adjusting significance thresholds for multiple comparisons.

Variation in weevil and plant phenologies. To compare magnitudes of year-to-year variation in *R. conicus* and *C. canescens* phenologies, we calculated coefficients of variation across years for proportion of *R. conicus* adults observed and proportion of *C. canescens* flower heads initiated by late May for each site with intensive count data in at least three years (Arapaho: RBO and West Foothills (3 yr); Niobrara: Inner and Outer Salzman (4 yr)). Because levels of floral herbivory can vary greatly over even small distances (Rodriguez et al. 1994; Traveset 1995), we also compared magnitudes of spatial variation in *R. conicus* and *C. canescens* phenologies by calculating coefficients of variation across sites for proportions of *R. conicus* adults observed and *C. canescens* flower heads initiated by late May. In each year, the sample included two Arapaho sites and two Niobrara sites. To evaluate whether

variation in weevil phenology or variation in thistle phenology was primarily responsible for generating variation in synchrony of their populations, we examined Pearson product moment correlations of both weevil and plant phenologies with the degree *R. conicus* preceded *C. canescens*.

We examined climate effects on variation in *C. canescens* reproductive phenology, 1990-2001, using multiple regression. Seasonal growing degree days and precipitation (spring: April-June, summer: July-September, autumn: October-December, winter: January-March) as well as days until last spring freeze and days until last spring hard freeze ($< 3.9^{\circ}\text{C}$) were calculated from daily climate data and used as explanatory variables. We standardized climate variables for each preserve by subtracting the mean of the variable and dividing by its standard deviation. To remove effects of variation in sampling date on our measures of plant phenology we used residuals from the regression of arcsine-transformed proportion of heads initiated by May on sampling date as our dependent variable in multiple regression models. We evaluated climate models according to over-all significance, mean square error, and inclusion of non-significant explanatory variables.

Effects of Phenological Synchrony on Egg Loads. To evaluate the effect of variation in phenological synchrony of *R. conicus* and *C. canescens* populations on variation in *R. conicus* egg loads on *C. canescens* among sites and years, we regressed log-transformed mean egg cases per flower head on the degree *R. conicus* preceded *C. canescens*, using simple, linear regression. With each site in each year, 1997-2000, as a separate unit of observation (Platte thistle is monocarpic, precluding re-sampling between years), we had 21 combinations of sites and years.

To evaluate the hypothesis that the magnitude of asynchrony, but not the identity of the species that was earlier (*R. conicus* or *C. canescens*), determined egg load (Louda 1998), we regressed log-transformed mean egg cases per flower head on the absolute value of the difference between the proportion of weevils observed and the mean proportion of *C. canescens* flower heads initiated, by late May. We refer to this absolute value as the “magnitude of asynchrony.”

Results

Variation in *R. conicus* Egg Loads

We found large variation among years and preserves in *R. conicus* egg load on *C. canescens*. At Niobrara in the year of highest use (1998), the mean number of eggs per flower head per plant (mean = 3.24) was 22X greater than egg load in the year of lowest use (2001, mean = 0.15), ($\chi^2 = 33.65$, df = 6, $p < 0.001$). At Arapaho, in the year of highest use (2001), on average 3.23 egg cases were observed per flower head whereas in 1996 we observed no egg cases ($\chi^2 = 38.45$, df = 6, $p < 0.001$). Mean egg load averaged across years at Arapaho Prairie (mean = 2.44) was 32.9% greater than at the Niobrara Preserve (mean = 1.84; $\chi^2 = 7.65$, df = 1, $p = 0.006$).

Variation in Insect and Plant Phenology

For the four years (1997 – 2000) when we counted *R. conicus* adults twice weekly, we found a significant year X preserve interaction effect on weevil phenology ($\chi^2 = 54.69$, df = 2, $p < 0.001$; Fig 1A,B). A significantly higher proportion of *R. conicus* adults had appeared by late May at Arapaho than at Niobrara in 1998 and 1999. Weevil phenology did not differ significantly among preserves in 1997 or 2000. For the 12 years (1990 – 2001)

when we sampled *C. canescens* plants, we also found a significant year X preserve interaction effect on *C. canescens* reproductive phenology (interaction: $F_{1,393} = 6.94$, $p = 0.001$). In 1991 and 1994, a significantly higher proportion of flower heads produced during the season were initiated by late May at Arapaho than at Niobrara. Platte thistle phenology did not differ between preserves in any other year (Fig. 1C,D).

The magnitudes of temporal and spatial variation in *R. conicus* phenology consistently exceeded those in *C. canescens* phenology between 1997 and 2000. At RBO (at Arapaho) and Inner and Outer Salzman (at Niobrara), coefficients of variation for year-to-year variation in *R. conicus* phenology were at least 4.0X as great as coefficients of year-to-year variation in *C. canescens* phenology (Table 3A). For West Foothills (at Arapaho), coefficients of variation for year-to-year variation in weevil and plant phenologies were equal. In each year, variation among sites in *R. conicus* phenology was greater than variation in *C. canescens* phenology (Table 3B). Coefficients of variation for spatial variation in *R. conicus* phenology were at least 10.0X the coefficients of variation for spatial variation in *C. canescens* phenology. The degree *R. conicus* preceded *C. canescens* was significantly correlated with *R. conicus* phenology ($r = 0.979$, $p < 0.0001$), whereas it was not correlated with *C. canescens* phenology ($r = -0.288$, $p = 0.208$).

None of the climate variables that we examined significantly explained variation in reproductive phenology of *C. canescens* populations. Our “best” regression model ($R^2 = 0.258$, $p = 0.0735$) included days to last spring hard freeze ($p = 0.0955$) and spring precipitation ($p = 0.1345$). Later hard freezes and lower spring precipitation were associated with lower proportions of flower heads initiated by late May.

Synchrony and Oviposition Intensity

Relative phenologies of *R. conicus* and *C. canescens* populations affected *R. conicus* use of *C. canescens*. The log-transformed number of egg cases per flower head per plant was significantly, positively related to the degree *R. conicus* preceded *C. canescens* ($F_{1,19} = 7.02$, $p = 0.016$, $R^2 = 0.269$). However, the magnitude of asynchrony, which ignores which species (weevil or plant) was earlier, was more effective in explaining temporal and spatial variation in *R. conicus* egg load on *C. canescens* ($F_{1,19} = 17.21$, $p < 0.001$, $R^2 = 0.472$). Egg loads were higher in years and sites where phenologies of *R. conicus* and *C. canescens* populations were more synchronized (Fig. 2). The fit of a quadratic model for the relationship between egg load and the degree *R. conicus* preceded *C. canescens* ($F_{1,19} = 17.31$, $p < 0.001$, $R^2 = 0.477$) was identical to the fit with magnitude of asynchrony as the explanatory variable (above).

Discussion

Phenological Synchrony and Interaction Strength

Our results show that floral herbivory by *R. conicus*, an exotic biocontrol weevil, on populations of Platte thistle (*C. canescens*), a native North American adopted host plant, varied significantly among sites and years. In general, the mechanisms that produce variation in rates of herbivory among host plant populations and among years in individual host plant populations are poorly understood. We found that temporal and spatial variation in the degree of phenological synchrony of *R. conicus* adult activity and *C. canescens* floral development strongly predicted variation in levels of herbivory between plant populations and years. Use of *C. canescens* flower heads by *R. conicus* decreased as the magnitude of asynchrony between weevil and plant populations increased. The identity of the species (weevil or plant) that was earlier was less important than the degree of synchrony.

Variation in *R. conicus* adult phenology was most important in determining degree of synchrony of herbivore and plant populations. Phenology of *C. canescens* floral development was much less variable in space and time than was *R. conicus* phenology. Even in the Sand Hills' continental climate, variation in temperature and precipitation did not strongly influence timing of *C. canescens* reproduction. The insensitivity of *C. canescens* reproductive phenology to climatic variation contrasts with our finding that *R. conicus* phenology is strongly influenced by winter and spring temperature and spring precipitation (Russell and Louda, in review). Large variation in *R. conicus* phenology in the new environment may reflect either exposure to novel, harsher and more variable environmental conditions or availability of alternative resources (Russell and Louda in review).

Large variation in rates of predispersal seed losses to floral herbivory is common among host plant populations (Jordano et al. 1990; Leimu et al. 2002) and among years (De Steven 1981; Evans et al. 1989). Such spatial and temporal variation in the intensity of herbivory can be critically important for dynamics of plant populations and communities, because it can determine landscape distributions and population sizes of host plants (Louda 1982, 1983; Rand 2002), especially when floral and predispersal seed herbivores severely limit recruitment (Louda and Potvin 1995a; Maron et al. 2002). Variation in herbivore population sizes or in the amount of plant resources available to flower-feeding insects often is hypothesized to explain differences in levels of damage among populations and years (De Steven 1981; Leimu et al. 2002). Qualitative comparisons of phenologies of some floral herbivores and their host plants have suggested that low levels of herbivory in some plant populations may coincide with large mis-matches in the timing of insect activity and plant floral development (Evans et al.

1989; Jordano et al. 1990). However, prior to this study, we found only one quantitative test of the hypothesis that intra-seasonal variation in the degree of phenological synchrony between inflorescence-feeding insect populations and their host plant populations drives variation in interaction strength (Rodriguez et al. 1994). Further tests of this seasonal phenological synchrony hypothesis for floral herbivore impact on interaction strength and plant population parameters are merited.

Measures of Seasonal Synchrony of Interacting Populations

Among the 18 studies we found that quantified within season phenological synchrony of consumer and resource populations, we distinguished three categories for the measures of synchrony that were used (Table 2). Seasonal phenological synchrony has been quantified as a comparison of: 1) number of days elapsed between the date of first appearance, median appearance or modal abundance of the consumer and resource populations; 2) thermal units accumulated between the date of first (or median) appearance or modal abundance of the consumer vs. the resource population; and, 3) difference between the proportion of the consumer population observed by a set date and the proportion of the resource population observed by the same date. Comparison of days elapsed to first (or median) appearance or modal abundance for both populations (category 1) was overwhelmingly the most common measure of seasonal phenological synchrony, used in 72% of the 18 studies.

Quantification of phenological synchrony as calendar or developmental time elapsed -between first appearances, median appearances or modal abundances of consumer and resource populations— allows evaluation of whether the phenological time difference between populations is large enough to affect either consumer or resource individuals. For example, if the rate of change in leaf quality after budbreak is known, measuring herbivore-

plant phenological synchrony in days elapsed between first appearances permits assessment of whether the first leaves to expand will be unpalatable when folivores appear. This measure of phenological synchrony, however, provides little information about the frequency distributions of phenologies among individuals in consumer and resource populations. Skewness and kurtosis of these distributions will strongly influence the proportion of each population affected by asynchrony and, hence, population growth. Comparing dates of modal abundances, rather than dates of first appearances, should improve prediction of effects on consumer and resource population dynamics because distribution modes likely represent the phenologies of larger proportions of individuals in the consumer and resource populations.

Measures of phenological synchrony that compare proportions of consumer and resource populations observed by a predetermined date indicate the proportion of individuals in a population that could be affected by any asynchrony. Such measures, however, often provide insufficient information to determine whether the temporal gap between consumers and resources is large enough to impact performance of asynchronous individuals. A measure of phenological synchrony that compares proportions of consumers and resources observed by a predetermined date, using a date based on the length of the phenological window during which resource individuals are vulnerable to consumers, though, could provide a robust predictor of the effect of variation in seasonal phenological synchrony on consumer and resource population dynamics.

Our measure of phenological synchrony for *R. conicus* and *C. canescens* populations, which was the proportion of the adult weevil population observed by late May minus the proportion of flower heads in the *C. canescens* population initiated by late May, is an example of the third category of measures of phenological synchrony above. Because our

goal was to explain variation in herbivory among host plant populations, we chose a measure of phenological synchrony that we expected would closely reflect variation in the proportion of host plant individuals and their floral resources were vulnerable to herbivory. The late May sampling date, which is related to the phenological window of weevil activity and individual flower head vulnerability in our system, was used to quantify phenological synchrony of *R. conicus* and *C. canescens* populations; our previous data indicated that late May was, on average, in the middle of the three week interval during which *R. conicus* oviposition on *C. canescens* was highest and flower head initiation was peaking. Therefore, we expected that the degree of synchrony in late May would be critical to determining weevil damage inflicted on Platte thistle populations.

Measures used to quantify reproductive synchrony of populations of masting plant species across years might be adapted to provide an additional measure of within-season synchrony of interacting populations. Reproductive synchrony of masting plant populations usually is quantified as the correlation coefficient for number of seeds produced by two populations over multiple years (Elkinton et al. 1996; Schnurr et al. 2002). Large, positive correlation coefficients indicate populations are synchronized in seed production, i.e., they mast in the same years and forego reproduction in the same years. Within a growing season, synchrony of consumer and resource populations might be quantified as the correlation coefficient for the number of consumer and resource individuals observed by time, i.e., each week or each month. Correlation coefficients of joint abundances of consumer and resource densities obtained for different populations could be used as explanatory variables in regression analyses to evaluate whether the degree of consumer-resource phenological synchrony affects reproduction or size of consumer and resource populations. A quantitative

comparison of the predictive abilities of different measures of phenological synchrony of consumer and resource populations has yet to be done, but would be a very useful next step.

Conclusion

Phenological mis-match between consumer and resource populations increasingly is recognized as a potentially important dimension of the intensity of interactions between consumer and resource species, one that is vulnerable to global climate change. However, quantitative evidence on the effect of such variation on the properties of resource populations is very limited. We found that variation in seasonal phenology of an exotic, invasive insect floral herbivore affected the phenological synchrony of its interaction with an adopted native North American host plant. Further, the degree of phenological synchrony of populations strongly influenced the magnitude of impact of the insect on the host plant. Degree of synchrony was important here, not which species was first. Such synchrony is likely to be important in determining outcomes and strengths of consumer-resource interactions more generally as well. Our findings suggest that prediction of the effects of invasive insect herbivores on indigenous plant populations will be improved by quantifying the synchrony and variation in insect and plant phenologies.

Acknowledgments. We thank the many students, colleagues, friends, and family members who contributed to the field data collection and discussions of thistle-insect interactions since 1990, as well as C. W. O'Brien for weevil identification and S. Kachman and A. M. Parkhurst for statistical advice. We are very grateful also to colleagues who aided our search for relevant phenological studies: A. K. Brody, V. K. Brown, D. E. Bunker, A. Buse, N. Cappucino, W. P. Carson, M. J. Crawley, R. F. Denno, E. W. Evans, S. N. Handel, D.W.

Inouye, R. Karban, R. J. Marquis, P.A. Morrow, R.B. Root, S. Y. Strauss and A. VanVliet.

The staffs of Cedar Point Biological Station and the staffs of the Niobrara Valley Preserve of The Nature Conservancy and Cedar Point Biological Station have provided continuing encouragement and valuable field support. Partial support for the long-term data collection on thistle-insect interactions was provided by grants from: University of Nebraska Research Council, Rodney Johnson and Katherine Ordway Stewardship Endowments of The Nature Conservancy, The National Science Foundation (DEB9221065, DEB9616210), and the USDA National Research Initiative (2001-35320-09882).

Literature Cited

- Alonso C, Herrera CM (2000) Seasonal variation in leaf characteristics and food selection by larval noctuids on an evergreen Mediterranean shrub. *Acta Oecologica* 21:257-265
- Angulo-Sandoval P, Aide TM (2000) Leaf phenology and leaf damage on saplings in the Luquillo Experimental Forest, Puerto Rico. *BioTropica* 32:415-422
- Augsburger CK (1981) Reproductive synchrony of a tropical shrub: Experimental studies on effects of pollinators and seed predators on *Hybanthus prunifolius* (Violaceae). *Ecology* 62:775-788
- Bertin RI (1982) The Ruby-throated Hummingbird and its major food plants: ranges, flowering phenology, and migration. *Can J Zoo* 60:210-219
- Buse A, Dury SJ, Woodburn RJ, Perrins CM, Good JEG (1999) Effects of elevated temperature on multi-species interactions: the case of Pedunculate Oak, Winter Moth and Tits. *Funct Ecol* 13:74-82
- Buse A, Good JEG (1996) Synchronization of larval emergence in winter moth (*Operophtera brumata* L.) and budburst in pedunculate oak (*Quercus robur* L.) under simulated climate change. *Ecol Entomol* 21:335-343
- Cappuccino N, Kareiva P (1985) Coping with a capricious environment: a population study of a rare pierid butterfly. *Ecology* 66:152-161
- Crawley MJ, Akhteruzzaman M (1988) Individual variation in the phenology of oak trees and its consequences for herbivorous insects. *Funct Ecol* 2:409-415

- De Steven D (1981) Abundance and survival of a seed-infesting weevil, *Pseudanthonomus hamamelidis* (Coleoptera: Curculionidae), on its variable fruiting host plant, witch-hazel (*Hamamelis virginiana*). *Ecol Entomol* 6:387-396
- Dewar RC, Watt AD (1992) Predicted changes in the synchrony of larval emergence and budburst under climatic warming. *Oecologia* 89:557-559
- Elkinton JS et al. (1996) Interactions between gypsy moths, white footed mice and acorns. *Ecology* 77:2332-2342
- Eriksson O (1995) Asynchronous flowering reduces seed predation in the perennial forest herb *Actaea spicata*. *Acta Oecologia* 16:195-203
- Evans EW, Smith CC, Gendron RP (1989) Timing of reproduction in a prairie legume: Seasonal impacts of insects consuming flowers and seeds. *Oecologia* 78:220-230
- Great Plains Flora Association (1986) *Flora of the Great Plains*. University Press of Kansas, Lawrence, KS
- Herrera CM, Jordano P, Guitian J, Traveset A (1998) Annual variability in seed production by woody plants and the masting concept: reassessment of principles and relationship to pollination and seed dispersal. *Amer Nat* 152:576-594
- Hill JK, Hodkinson ID (1992) Effects of temperature on phenological synchrony and altitudinal distribution of jumping plant lice (Hemiptera: Psylloidea) on dwarf willow (*Salix lapponum*) in Norway. *Ecol Entomol* 20:237-244
- Hodkinson ID (1997) Progressive restriction of host plant exploitation along a climatic gradient: the willow psyllid *Cacopsylla groenlandica* in Greenland. *Ecol Entomol* 22:47-54

- Hodkinson ID, Bird JM, Hill JK, Baxter P (2001) Host plant growth characteristics as determinants of abundance and phenology in jumping plant-lice on downy willow. *Ecol Entomol* 26:371-387
- Hunter AF, Elkinton J (2000) Effects of synchrony with host plant on populations of a spring-feeding Lepidopteran. *Ecology* 81:1248-1261
- Hunter MD (1992) A variable insect-plant interaction: the relationship between tree budburst phenology and population levels of insect herbivores among trees. *Ecol Entomol* 16:91-95
- Inouye DW, Barr B, Armitage KB, Inouye BD (2000) Climate change is affecting altitudinal migrants and hibernating species. *PNAS* 97:1630-1633
- Jordano D, Fernandez Haeger J, Rodriguez J (1990) The effect of seed predation by *Tomares ballus* (Lepidoptera: Lycaenidae) on *Astragalus lusitanicus* (Fabaceae): Determinants of differences among patches. *Oikos* 57:250-256
- Kaitaniemi P, Ruohomaki K, Haukioja E (1997) Consequences of defoliation on phenological interaction between *Epirrita autumnata* and its host plant, Mountain Birch. *Funct Ecol* 11:199-208
- Kennedy AA, McCullough DG (2002) Phenology of the Larger European Pine Shoot Beetle *Tomicus piniperda* (L.) (Coleoptera: Scolytidae) in relation to native bark beetles and natural enemies in pine stands. *Environ Entomol* 31:261-272
- Kerslake JE, Hartley SE (1997) Phenology of winter moth feeding on common heather: effects of source population and experimental manipulation of hatch dates. *J Anim Ecol* 66:375-385

- Lamp WO (1980) Predispersal seed predation of the Platte Thistle and its effect on seed production. Ph.D. Thesis. University of Nebraska, Lincoln, p 173
- Lawrence RK, Mattson WJ, Haack RA (1997) White spruce and the spruce budworm: defining the phenological window of susceptibility. *Can Entomol* 129:291-318
- Leimu R, Syrjanen K, Ehrlén J, Lehtilä KP (2002) Pre-dispersal seed predation in *Primula veris*: among-population variation in damage intensity and selection on flower number. *Oecologia* 133:510-516
- Louda SM (1998) Population growth of *Rhinocyllus conicus* (Coleoptera: Curculionidae) on two species of native thistles in prairie. *Environ Entomol* 27:834-841
- Louda SM (1999) Negative ecological effects of the musk thistle biocontrol agent, *Rhinocyllus conicus* Fröl. In: Follet PA, Duan JJ (eds) Nontarget effects of biological control. Kluwer Academic Publishers, Boston, MA, pp 215-243
- Louda SM, Arnett AE (2000) Predicting non-target ecological effects of biological control agents: Evidence from *Rhinocyllus conicus*. In: Spencer NR (ed) Proc 10th Intl Sym Biol Cont Weeds. Montana State University, Bozeman, MT, pp 551-567
- Louda SM, Kendall D, Connor J, Simberloff D (1997) Ecological effects of an insect introduced for the biological control of weeds. *Science* 277:1088-1090
- Louda SM, Potvin MA (1995a) Effect of inflorescence-feeding insects in the demography and lifetime fitness of a native plant. *Ecology* 76:229-245
- Louda SM, Potvin MA (1995b) Effect of inflorescence-feeding insects on the demography and lifetime fitness of a native plant. *Ecology* 76:229-245
- Maron JL, Combs JK, Louda SM (2002) Insect herbivory on thistles: Congruent interaction strengths in coastal vs. continental dunes. *Ecology* 83:3382-3392

- Martel J, Hanhimaki S, Kause A, Haukioja E (2001) Diversity of birch sawfly responses to seasonally atypical diets. *Entomol Exp Appl* 100:301-309
- Mopper S, Simberloff D (1995) Differential herbivory in an oak population: The role of plant phenology and insect performance. *Ecology* 76:1233-1241
- Ollerton J, Lack A (1998) Relationships between flowering phenology, plant size and reproductive success in *Lotus corniculatus* (Fabaceae). *Plant Ecol.* 139:35-47
- Peterson MA (1997) Host plant phenology and butterfly dispersal: causes and consequences of uphill movement. *Ecology* 78:167-180
- Pilson D (2000) Herbivory and natural selection on flowering phenology in wild sunflower, *Helianthus annuus*. *Oecologia* 122:72-82
- Quiring DT, McKinnon ML (1999) Why does early-season herbivory affect subsequent budburst? *Ecology* 80:1724-1735
- Rathcke B, Lacey EP (1985) Phenological patterns of terrestrial plants. *Ann Rev Ecol Syst* 16:179-214
- Rodriguez J, Jordano D, Fernandez Haeger J (1994) Spatial heterogeneity in a butterfly-host plant interaction. *J Anim Ecol* 63:31-38
- Schnurr JL, Ostfeld RS, Canham CD (2002) Direct and indirect effects of masting on rodent populations and tree seed survival. *Oikos* 96:402-410
- Sokal RR, Rohlf FJ (1995) *Biometry*, 3rd edn. W.H. Freeman and Company, New York
- Stenseth NC, Mysterud A (2002) Climate, changing phenology, and other life history traits: Nonlinearity and match-mismatch to the environment. *PNAS* 99:13379-13381

- Surles WW, Kok LT (1977) Ovipositional preference and synchronization of *Rhinocyllus conicus* with *Carduus nutans* and *C. acanthoides*. *Environ Entomol* 6:222-224
- Thireau JC, Regniere J (1995) Development, reproduction, voltinism and host synchrony of *Meteorus trachynotus* with its host *Choristoneura fumiferana* and *C. rosaceana*. *Entomol Exp Appl* 76:67-82
- Tikkanen O, Carr TG, Roininen H (1999) Factors influencing the distribution of a generalist spring-feeding moth, *Operophtera brumata* (Lepidoptera: Geometridae), on host plants. *Environ Entomol* 28:461-469
- Traveset A (1995) Spatio-temporal variation in pre-dispersal reproductive losses of a Mediterranean shrub, *Euphorbia dendroides* L. *Oecologia* 103:118-126
- Van Noordwijk AJ, McCleery RH, Perrins CM (1995) Selection for the timing of Great Tit breeding in relations to caterpillar growth and temperature. *J Anim Ecol* 64:451-458
- Visser ME, Holleman LJM (2001) Warmer springs disrupt the synchrony of oak and winter moth phenology. *Proc R Soc Lond Ser B* 268:289-294
- Volney WJA, Cerezke HF (1992) The phenology of white spruce and the spruce budworm in northern Alberta. *Can J For Res* 22:198-205
- Watt AD, McFarlane AM (1991) Winter moth on Sitka spruce: synchrony of egg hatch and budburst and its effect on larval survival. *Ecol Entomol* 16:387-390
- Zwölfer H, Harris P (1984) Biology and host specificity of *Rhinocyllus conicus* (Froel.) (Col., Curculionidae), a successful agent for biocontrol of the thistle, *Carduus nutans* L. *Z Angew Entomol* 97:36-62

Figure 1. *Rhinocyllus conicus* adult phenology, *Cirsium canescens* phenology, and phenological synchrony of *R. conicus* with *C. canescens* flower heads at Niobrara Valley Preserve (A,C,E) and at Arapaho Prairie Preserve (B,D,F) 1997 – 2000: A, B = within-season temporal distributions of *R. conicus* adults, showing running means of the number of *R. conicus* adults standardized per 20 thistle plants per preserve; C, D = mean (S.D.) percent of the seasonal total *C. canescens* flower heads per plant initiated by late May; E, F = degree *R. conicus* preceded *C. canescens* (% *R. conicus* adults observed - % *C. canescens* heads initiated, by late May); larger absolute values indicate greater asynchrony than smaller absolute values. In C-E, years are ranked by earliest to latest *R. conicus* phenology. For each preserve, “a” = year of earliest (fastest development) *R. conicus* phenology, while “d” = year of latest (slowest development) *R. conicus* phenology.

Figure 2. A) *Rhinocyllus conicus* oviposition on *Cirsium canescens* flower heads per plant as a function of the magnitude of asynchrony of phenology of the *R. conicus* and *C. canescens* populations. Magnitude of asynchrony is quantified as the absolute value of the difference between the proportion of *R. conicus* observed – the proportion of *C. canescens* heads initiated by late May. Larger absolute values indicate greater asynchrony than smaller absolute values. B) Oviposition on *C. canescens* as a function of the degree *R. conicus* preceded *C. canescens*. Positive values indicate *R. conicus* earlier than *C. canescens*. Negative values indicate *R. conicus* later than *C. canescens*. Solid circles represent Niobrara sites, and open circles represent Arapaho sites. In both panels, site means for number of egg cases per flower head per individual plant ranged from 5.76 egg cases / head (Mahoney site (NVP) in 1998) to 0.38 egg cases / head (Inner Salzman site (NVP) in 1999).

Table 1. Questions generally asked about the ecological and evolutionary effects of seasonal phenology of interacting phytophagous insects and host plants (1993 – 2002), with example studies. Phenology is defined as seasonal timing within year (Rathcke and Lacey 1985), so excludes masting.

1. Does herbivory affect leaf or reproductive phenology of an individual plant in future growing seasons? (Kaitaniemi et al. 1997; Quiring and McKinnon 1999);
 2. Does individual plant phenology affect the amount of herbivore damage to it? (Angulo-Sandoval and Aide 2000; Mopper and Simberloff 1995);
 3. Do herbivores affect evolution of individual plant leaf or reproductive phenology? (Ollerton and Lack 1998; Pilson 2000);
 4. Does plant species phenology determine host species use by insects? (Hodkinson 1997; Tikkanen et al. 1999);
 5. Does year-to-year variation in the synchrony of insect and plant populations affect insect population size, especially outbreaks? (Hunter and Elkinton 2000; Kerslake and Hartley 1997);
 6. Does variation in plant phenology explain spatial structure of phytophagous insect abundances, such as among individual plants? (Hodkinson et al. 2001; Rodriguez et al. 1994);
 7. Does host plant phenology explain seasonal variation in herbivorous insect abundance? (Alonso and Herrera 2000) or migratory patterns (Peterson 1997);
 8. Does insect phenology relative to plant phenology determine response to novel plant tissue quality? (Martel et al. 2001).
-

Table 2. Studies that quantify magnitude and variation in seasonal phenological synchrony of field populations of consumer and resource species. Pollination studies are included because pollinators often consume nectar or pollen. Measures of seasonal phenological synchrony were: 1) “time difference” = calendar days elapsed between first (or median) appearance or modal abundance of consumer and resource; 2) “thermal difference” = difference in thermal units accumulated between first (or median) appearance or modal abundance of consumer and resource; and, 3) “Proportion appeared” = difference between proportion of consumer population present and proportion of resource population available on a fixed date. For both “time difference” and “thermal difference,” three basic distribution parameters have been used: first appearance, median appearance, modal abundance; these are indicated parenthetically.

Species (consumer; resource)	Phenological synchrony measured as:	Phenological synchrony used as:	Population effects quantified for:	Source
Herbivory				
<i>Cacopsylla brunneipennis, C. palmeni; Salix lapponum</i>	Thermal difference (mean)	Independent variable	Consumer	(Hill and Hodkinson 1992)
<i>Choristoneura fumiferana; Picea glauca</i>	Time difference (first)	Independent variable	Consumer	(Lawrence et al. 1997)

Species (consumer; resource)	Phenological synchrony measured as:	Phenological synchrony used as:	Population effects quantified for:	Source
Herbivory (cont.)				
<i>Cyaniris semiargus</i> ; <i>Armeria velutina</i>	Time difference (mode)	Independent variable	Consumer	(Rodriguez et al. 1994)
<i>Lymantria dispar</i> ; <i>Quercus velutina</i> , <i>Q.</i> <i>alba</i>	Time difference (first)	Independent variable	Consumer	(Hunter and Elkinton 2000)
<i>Operophtera</i> <i>brumata</i> ; <i>Picea</i> <i>sitchensis</i>	Proportion appeared	Independent variable	Consumer	(Watt and McFarlane 1991)
<i>Choristoneura</i> <i>fumiferana</i> ; <i>Picea</i> <i>glauca</i>	Thermal difference (multiple distribution parameters)	Dependent variable	Neither	(Volney and Cerezke 1992)

Species (consumer; resource)	Phenological synchrony measured as:	Phenological synchrony used as:	Population effects quantified for:	Source
---------------------------------	---	---------------------------------------	--	--------

Herbivory (cont.)

<i>Marmota flaviventris</i> ; plant community	Time difference (first)	Dependent variable	Neither	(Inouye et al. 2000)
<i>Operophtera brumata</i> ; <i>Quercus robur</i>	Time difference (median, mean)	Dependent variable	Neither	(Visser and Holleman 2001)
<i>Operophtera brumata</i> ; <i>Quercus robur</i>	Time difference (median, mean)	Dependent variable	Neither	(Buse and Good 1996)
<i>Operophtera brumata</i> ; <i>Quercus robur</i>	Time difference (mean)	Dependent variable	Neither	(Buse et al. 1999)

Species (consumer; resource)	Phenological synchrony measured as:	Phenological synchrony used as:	Population effects quantified for:	Source
---------------------------------	---	---------------------------------------	--	--------

Herbivory (cont.)

<i>Operophtera brumata; Picea sitchensis</i>	Time difference (median)	Dependent variable	Neither	(Dewar and Watt 1992)
<i>Pieris virginiensis; Dentaria diphylla</i>	Proportion appeared	Dependent variable	Neither	(Cappuccino and Kareiva 1985)
<i>Rhinocyllus conicus; Carduus nutans, C. acanthoides</i>	Proportion appeared	Dependent variable	Neither	(Surles and Kok 1977)

Pollination

<i>Archilochus colubris; Impatiens biflora, Aesculus pavia, Aquilegia canadensis, Castilleja coccinea</i>	Time difference (mode)	Dependent variable	Neither	(Bertin 1982)
---	------------------------------	-----------------------	---------	---------------

Species (consumer; resource)	Phenological synchrony measured as:	Phenological synchrony used as:	Population effects quantified for:	Source
---------------------------------	---	---------------------------------------	--	--------

Predation and Parasitism

<i>Parus major</i> ; <i>Operophtera brumata</i>	Time difference (median)	Independent variable	Consumer	(Van Noordwijk et al. 1995)
<i>Parus major</i> ; <i>Operophtera brumata</i>	Time difference (mode)	Independent & dependent variables	Consumer	(Buse et al. 1999)
<i>Cucujus clavipes</i> , <i>Enoclerus nigripes</i> , <i>Platysoma</i> <i>cylindrical</i> , <i>P.</i> <i>parallellum</i> , <i>Thanasimus dubius</i> , <i>Tomicus piniperda</i> ; <i>Corticeus parallelus</i>	Time difference (first, mode)	Dependent variable	Neither	(Kennedy and McCullough 2002)

Species (consumer; resource)	Phenological synchrony measured as:	Phenological synchrony used as:	Population effects quantified for:	Source
---------------------------------	---	---------------------------------------	--	--------

Predation and Parasitism (cont.)

<i>Turdus migratorius</i> ; insect community	Time difference (first)	Dependent variable	Neither	(Inouye et al. 2000)
<i>Meteorus trachynotus</i> ; <i>Choristoneura fumiferana</i> , <i>C. rosaceana</i>	Time difference (mode)	Dependent variable	Neither	(Thireau and Regniere 1995)

Table 3. Coefficients of variation for temporal (A) and spatial (B) variation in phenologies of adult activity of *R. conicus* populations and flower head development of *C. canescens* populations.

A) Variation among years for individual sites

Site	Preserve	Coefficient of variation, <i>R. conicus</i> phenology	Coefficient of variation, <i>C. canescens</i> phenology
Roses Blowout	Arapaho	187.58	17.26
West Foothills	Arapaho	15.21	15.42
Inner Salzman	Niobrara	54.68	13.67
Outer Salzman	Niobrara	149.07	11.29

B) Variation among sites for individual years

Year	Coefficient of Variation, <i>R. conicus</i> phenology	Coefficient of Variation, <i>C. canescens</i> phenology
1997	214.08	12.18
1998	123.08	11.83
1999	117.25	7.76
2000	78.05	6.36



